

ORGANIC MATTER EFFECT ON *Glomus intraradices* IN BEANS (*Phaseolus vulgaris* L.) GROWTH CULTIVATED IN SOILS WITH TWO SOURCES OF WATER UNDER GREENHOUSE CONDITIONS

Gardezi, A. K.¹, Márquez-Berber, S. R.², Figueroa-Sandoval, B.¹, Larqué-Saavedra U.³, Almaguer-Vargas, G.², Escalona-Maurice M. J.⁴.

¹ Colegio de Postgraduados. Instituto de Recursos Naturales, Programa de Hidrociencias. Montecillo, Texcoco, Estado de México. Tel (595) 9520200 ext. 1168, email: kabdul@colpos.mx. ² Universidad Autónoma Chapingo. Departamento de Fitotecnia. ³ Universidad Autónoma Metropolitana-Azcapotzalco. Área de Estadística e Investigación de Operaciones. ⁴ Colegio de Posgraduados. Campus Montecillo. Desarrollo Rural.

SUMMARY

The objective of this research was to evaluate the effect of organic matter on the association with *Glomus intraradices* and soil contamination on beans (*Phaseolus vulgaris* L.). The study was done under greenhouse conditions at the Montecillo Campus of the Postgraduate College, Mexico. Two soils were used, one irrigated with sewage water and the other one with clean water from a well. Half of the plants were inoculated with *Glomus intraradices*. Vermicompost was used as a source of organic matter. There were highly significant increases ($p \leq 0.05$) in all the variables recorded due to the application of organic matter, and to the inoculation with *Glomus intraradices*. The irrigation source of the soils used in this experiment only had a significant effect ($p \leq 0.05$) on pod number and nitrogen fixation. The best growth and grain yield occurred with inoculated plants and supplementary organic matter.

Key words: Soil contamination, vermicompost, sewage water, arbuscular endomycorrhiza, edible legumes.

INTRODUCTION

Common bean is the most important legume in Mexico. More than 1.5 million hectares were planted in 2011 in this country [24]. It is cultivated mainly in areas where water is scarce. Sewage water has been used for its irrigation. Studies on its effect on plant growth and yield are needed. Farmers usually apply chemical fertilizer. Its cost has been increasing dramatically [18]. Organic matter sometimes is applied as fertilizer. However, their consequences on beneficial microorganisms as endomycorrhiza need to be understood. *Glomus intraradices* has been used to improve plant growth under different conditions included contaminated soils. Several researchers consider that kind of fungi as the most important organisms on earth interacting in agro environments. More than 80% of all terrestrial plants, among them most of the horticultural and crop plants have a symbiotic relationship with these fungi. The stimulation of plant growth can be attributed mainly to the

improvement of phosphorus nutrition [1, 9, 12, 17, 23]. *Glomus intraradices* has increased bean yield 36% [19]. Novella *et al.* [21] had reported augmented corn and bean yield when they were cultivated together and were inoculated with a combination of *Rhizobium* and mycorrhiza.

Glomus mosseae amplified shoot growth four times [27]. Mycorrhizal development was better under no tillage conditions than using conventional one in an oat-wheat rotation [6]. Mineral nitrogen improves biomass and root growth when mycorrhiza is present [26].

The objective of this study was to investigate the effect of organic matter associated with *Glomus intraradices*, on the growth and yield of beans, in two soils with different water source under greenhouse conditions.

MATERIALS AND METHODS

The study was done under greenhouse conditions at the Postgraduate College, Montecillo Campus, State of Mexico, in the spring and summer of 2012. Two soils, from Tocuila, Texcoco, Mexico, were used. One was irrigated with sewage water and the other one with clean water from a well. Their characteristics are shown in Table 1.

The seeds were sterilized with 1% sodium hypochlorite during 4 minutes, and hydrated on filter paper in petri dishes for 48 hours. The seeds were sown in polyethylene bags that had been filled with three kg of two soil types. The treatments were: planting in soil irrigated with sewage water and the other one with clean water. Both soils were collected at three depths 0-5, 5-10, and 10-40 cm from plots of one hectare [5].

The soil organic matter was determined using the Walkey and Black method. For phosphorus, Olsen was utilized. Interchangeable bases were measured with ammonium acetate 1 Normal ($\text{CH}_3\text{COONH}_4$) pH 7, and micronutrients with DTPA (diethylene-triamine-pentaacetic acid).

Bayomex bean genotype was planted. The seed was provided by the National Institute for Agriculture, Livestock and Forestry Research (INIFAP).

The inoculation was done during the planting, mixing 5 g of sand with sorghum roots with 78 % colonization of *Glomus intraradices* and 1050 spores per 100 g of inert

material. Two levels of *Glomus* were applied, with and without *Glomus*.

Organic matter was applied as a vermicompost. It was prepared using 60 kg of bovine manure, 25 kg of melon waste, and 15 kg of wheat straw. The mixture was subjected four months to the action of earth worms. Four doses were applied. In every bag of three kg, 0, 28.86 g, 57.7 g, and 86.46 g of vermicompost were mixed. They are equivalent to 0, 25, 50, and 75 t ha⁻¹ of organic matter.

Table 1 Soil analysis for the two types, one irrigated with sewage water and the other with clean one.

	pH	EC	OM	TN	NO ₃	P	CEC
Soil sample	1:2	dm sec ⁻¹	%	%	mg kg ⁻¹	mg kg ⁻¹	C mol (+) kg ⁻¹
Soil depth 0-5	7.44	349	2.5	0.098	18	15	19
Residual water							
Soil depth 5-10	7.37	454	2.48	0.096	17	14	18
Residual water							
Soil depth 10-40	7.44	475	2.45	0.094	15	11	15
Residual water							
Soil depth 0-5	7.52	314	2.49	0.097	17	14	18
Clean water							
Soil depth 5-10	7.75	332	2.47	0.095	16	13	17
Clean water							
Soil depth 10-40	7.85	384	2.43	0.092	13	10	13
Clean water							
Soil sample	Ca	Mg	K	Na	Fe	Zn	Cu
	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹
Soil depth 0-5	1250	59	1180	400	5	3	0.5
Residual water							
Soil depth 5-10	1210	56	1220	640	5	3	0.5
Residual water							
Soil depth 10-40	1245	59	1400	800	4	3	0.4
Residual water							
Soil depth 0-5	1240	58	1120	360	5	3	0.5
Clean water							
Soil depth 5-10	1243	60	1110	640	3	3	0.4
Clean water							
Soil depth 10-40	1200	53	1140	720	4	2	0.4
Clean water							

Key: pH= Hydrogen potential, EC=Electrical conductivity, OM= Organic matter, TN= Total nitrogen, NO₃= Nitric nitrogen, CEC= Cation Exchange Capacity.

The variables evaluated were: plant height (PH, cm), leaf area (LA, cm²), pod number (PN), grain dry weight (GDW, g), root length (RL, cm), root volume (RV cm³), root dry weight (RDW, g), pod dry weight (PDW, g), grain number (GN) biomass dry weight (BDW, g), white nodule number (WNN), red nodule number (RNN), and total nodule number (TNN).

A factorial arrangement with 16 treatments (4x2x2) was used with a completely randomized block

design with three replications. An analysis of variance for all variables registered was done and a Tukey mean comparison test for the significant variables.

RESULTS AND DISCUSSION:

The soil texture was sandy loam. The pH was alkaline. It was higher in the soils irrigated with clean water. The difference was greater in the 10-40 cm depth. The soil with clean water had a pH of 7.85, and that with sewage water only 7.44 (Table 1). The electrical conductivity (EC), organic matter (OM), total nitrogen (TN), nitric nitrogen (NO₃), phosphorous (P), Cation Exchange Capacity (CEC), calcium (Ca), potassium (K), sodium (Na), iron (Fe), zinc (Zn), and copper (Cu) quantities are higher in the soils that were irrigated with sewage water. In both soils, total nitrogen (TN), and the nitric nitrogen (NO₃) quantities were low. They distribution were higher in the 5-10 cm layer. No ammoniacal nitrogen (NH₄) was found in the two soils. It explains why total nitrogen is only slightly higher on the soils that were irrigated with sewage water that have higher organic matter [20].

The levels of Cu are below the threshold for considering them as contaminants [5]. No Cr and Ni traces were found.

There were highly significant differences (p≤0.01) in all the variables recorded due to the application of organic matter, and to the inoculation with *Glomus intraradices*. The pod number, and the white, red, and total nodule number were significantly affected (p≤0.05) for the contamination from the soil with sewage water. Their higher content of nitric and total nitrogen could reduce the nodule number. It is well known the antagonistic effect between nitrogen content in soils and nitrogen fixation [7]. The lack of effect on growth and yield could be explained by the low N fixation [8, 22].

A positive effect of inoculation with *Glomus intraradices* was found. The highly significant differences (p≤0.01) among treatments for all the variables recorded generated a beneficial effect on plant growth due to an improvement in the absorption of mineral nutrients required by the plants [2, 3]. This behavior was similar to that found by Gardezi *et al.* [8, 9, 12, 13 and 14].

Yield, root and shoot growth from plants with mycorrhiza were superior to those without inoculation (Table 2). The treatments with mycorrhiza were 54% taller, had 25% more leaf area (Figure 2), 48% heavier shoot dry weight, 47% longer roots, 56% greater root volume, and 48% more root dry weight¹ (Figure 3). This is an indication of a positive effect of mycorrhiza on

¹ Increase percentages are referred to the values found in bean plants inoculated with mycorrhiza compared to those without inoculation.

plant growth originated by better mineral nutrient absorption required by the plant [1, 2]. Gardezi *et al.* [10, 11, 12] also found this beneficial effect in *Leucaena leucocephala* associated with endomycorrhiza and with *Rhizobium*. Positive responses to the inoculation with mycorrhiza were also found in a number of species [24], including beans [2].

The inoculation with *Glomus intraradices* improved root and shoot growth. It also had a beneficial effect on the biological nitrogen fixation, and a superior absorption of nutrients [12]. Thus, it contributed to higher yield in beans, coinciding with other studies [2]. Inoculated plants had 38% heavier pods and yielded 116% more grain (Table 2, and Figure 1).

Table 2. Honest significant difference of the effect of *Glomus intraradices* on common bean (*Phaseolus vulgaris*).

Glomus intraradices	Plant height (cm)	Dry weight aerial part (g)	Leaf area (cm ²)	Grain number (ln)	Grain dry weight (g)	Pod dry weight (g)	Pod number (ln)
Inoculated	124.542a	24.292a	448.79a	2.9806a	10.958a	3.0000a	2.1849a
No inoculated	100.667b	16.375b	357.92b	2.3845b	5.083b	2.1667b	1.8475b

Glomus intraradices	Root length (cm)	Root volume (cm ³)	Dry weight root (g)	White nodule number (ln)	Red nodule number (ln)	Total nodule number (ln)
Inoculated	28.708a	9.7917a	3.0833a	1.9818a	2.6397a	3.0620a
No inoculated	19.500b	6.2500b	2.0833b	1.2320b	1.9258b	2.2557b

ln=transformed to natural log. Means with the same letter in each column are not significantly different (Tukey $\alpha=0.05$)




Figure 1. Effect of inoculation with *Glomus intraradices* and different doses of organic matter (vermicompost) on soils with two types of irrigation in grain dry weight of common bean (*Phaseolus vulgaris*).

Key: Glumus=0: Noninoculated, Glumus=1: Inoculated with *Glomus intraradices*. The vertical lines indicate standard error.

Plant growth was affected by the organic matter application (Table 3). It provided significantly higher ($p\leq 0.01$) plant height in all the treatments compared with the control. Aryal *et al* [3] found similar results. Only the higher quantities gave heavier dry weight of the aerial part (Figure 4). A similar situation was found in the leaf area (Figure 2).

Table 3. Honest significant difference of the effect of organic matter (vermicompost) on common bean (*Phaseolus vulgaris*).

Organic matter t*ha ⁻¹	Plant height (cm)	Dry weight aerial part (g)	Leaf area (cm ²)	Grain number (ln)	Grain dry weight (g)	Pod dry weight (g)	Pod number (ln)
0	90.167b	15.333b	361.67b	2.289b	4.833b	2.083b	1.824b
25	115.000a	19.833ab	402.50ab	2.511ab	7.917ab	2.667a	1.949ab
50	123.417a	22.917a	415.00a	2.963a	6.083b	2.750a	2.131a
75	121.833a	23.250a	434.25a	2.968a	13.250a	2.833a	2.161a

Organic matter t*ha ⁻¹	Root length (cm)	Root volume (cm ³)	Dry weight root (g)	White nodule number (ln)	Red nodule number (ln)	Total nodule number (ln)
0	19.583b	5.917b	2.000b	1.125b	1.744b	2.070b
25	24.833ab	7.667ab	2.750a	1.637ab	2.564a	2.919a
50	25.917a	9.083a	3.000a	1.682ab	2.337ab	2.692a
75	26.083a	9.417a	2.583a	1.983a	2.487a	2.954a

ln=transformed to natural log. Means with the same letter in each column are not significantly different (Tukey $\alpha=0.05$)




Figure 2. Effect of inoculation with *Glomus intraradices* on soils with two types of irrigation on leaf area of three cultivars common bean (*Phaseolus vulgaris*).

Key: Glumus=0: Noninoculated, Glumus=1: Inoculated with *Glomus intraradices*. The vertical lines indicate standard error.

Root length and root volume (Figure 5) were also significantly greater ($p\leq 0.05$) only with the two higher applications of vermicompost. However, all the doses of organic matter gave heavier roots.

Organic matter also promoted nitrogen fixation. All the vermicompost applications had a significantly higher total nodule number. However, only the elevated dose was related with a greater white and red nodule number.




Figure 3. Effect of inoculation with *Glomus intraradices* and organic matter application (vermicompost) in soils with two types of irrigation on pod number of common bean (*Phaseolus vulgaris*).

Key: Glumus=0: Noninoculated, Glumus=1: Inoculated with *Glomus intraradices*. The vertical lines indicate standard error.




Figure 4. Effect of inoculation with *Glomus intraradices* and organic matter application (vermicompost) in soils with two types of irrigation on plant height of common bean (*Phaseolus vulgaris*).

Key: Glumus=0: Noninoculated, Glumus=1: Inoculated with *Glomus intraradices*. The vertical lines indicate standard error.

In an analogous way with mycorrhizal inoculation, the organic matter promoted better root and shoot growth. Therefore, as a result, photosynthetic production increased. Pod number was higher with

organic matter (Figure 3). Grain yield was 174% enhanced with the highest dose of organic matter compared to the control (Figure 1, Table 3).

Thus, the poorest growth and grain yield occurred with uninoculated that lacked supplementary organic matter. The irrigation source of the soils used in this experiment only had a significant effect ($p \leq 0.05$) on pod number and nitrogen fixation. The higher content of organic matter and nitrogen found in the soils watered with sewage water generated a greater pod number. However, this effect was not present in grain yield (Table 4).

Table 4. Honest significant difference of the effect of the irrigation source on common bean (*Phaseolus vulgaris*).

Irrigation Source	Plant height (cm)	Dry weigh aerial part (g)	Leaf area (cm ²)	Grain number (ln)	Grain dry weight (g)	Pod dry weight (g)	Pod number (ln)
Clean water	109.17a	19.38a	400.88a	2.64a	8.46a	2.58a	1.93b
Sewage water	115.29a	21.29a	405.83a	2.74a	7.58a	2.58a	2.10a
Irrigation Source	Root lenght (cm)	Root volume (cm ³)	Dry weigh root (g)	White nodule number (ln)	Red nodule number (ln)	Total nodule number (ln)	
Clean water	23.46a	7.50a	2.46a	2.31a	2.71a	3.23a	
Sewage water	24.75a	8.54a	2.71a	0.91b	1.85b	2.09b	

ln=transformed to natural log. Means with the same letter in each column are not significantly different (Tukey $\alpha = 0.05$)




Figure 5. Effect of inoculation with *Glomus intraradices* and organic matter application (vermicompost) in soils with two types of irrigation on root volume of common bean (*Phaseolus vulgaris*).

Key: Glumus=0: Noninoculated, Glumus=1: Inoculated with *Glomus intraradices*. The vertical lines indicate standard error.

The nitrogen fixation, measured as the nodule number (white, red, and total) was higher in the soils irrigated with clean water (Table 4). A plausible explanation for this finding is the antagonistic effect between nitrogen content in soils and nitrogen fixation [7]. The soils watered with sewage water had higher total and nitric nitrogen (Table 1). Gardezi *et al.* [15, 16] found similar results in previous experiments.

CONCLUSIONS

Mycorrhizal inoculation and nitrogen fixation provided higher bean root and shoot growth and therefore, better yields. Previous evidence with legumes showed that they have benefited with this symbiosis because the treatments with this fungus produce the highest values for all evaluated variables. The application of organic matter, as vermicompost, improved plant growth, and grain yield. The contamination of the soil by sewage water did not affect plant growth or yield. It only affected nitrogen fixation.

ACKNOWLEDGMENTS

The authors want to thank Dr. Juan Antonio Villanueva, Head of Research, Montecillo Campus, Colegio de Postgraduados, for his valuable help for the development of this project.

LITERATURE CITED

[1] Alarcón, A. 2008. Los hongos micorrízicos arbusculares como biotecnología en la propagación y manejo de plantas en viveros. *Agroproductividad* 1(1): 19-23

[2] Aryal, U. K., H. L. Xu, and M. Fujita. 2003. Rhizobia and AM fungal inoculation improve growth and nutrient uptake of bean plants under organic fertilization. *J Sustainable Agric.* 21(3): 27-39.

[3] Aryal, U. K., S. K. Shah, H. L. Xu, M. Fujita. 2006. Growth, nodulation and mycorrhizal colonization in bean plants improved by rhizobial inoculation with organic and chemical fertilization. *J Sustainable Agric.* 29(2): 71-83.

[4] Bermúdez, M., L. Castro, y R. Araya V. 1995. Efecto de la aplicación de micorrizas y roca fosfórica sobre el crecimiento y la adquisición de nutrientes de plantas de frijol en un ultisol de Costa Rica. Mejoramiento y manejo agronómico del frijol común (*Phaseolus vulgaris* L.) para adaptación en suelos de bajo fósforo. *En* Beck, D.; comp. y ed. Memoria del Taller Internacional sobre

Bajo Fósforo en el Cultivo de Frijol. San José (Costa Rica). 13-15 Nov 1995. p. 95-103.

[5] Castellanos, J. Z., J. X. Uvalle B. y A. Aguilar S. 2000. Manual de interpretación de suelos y aguas. Colección INCAPA. México. 226 p.

Castillo *et al.*, 2006

[6] Castillo, C. G.; Rubio, R.; Rouanet, J. L.; Borie, F. 2006. Early effects of tillage and crop rotation on arbuscular mycorrhizal fungal propagules in an Ultisol. *Biology and Fertility of Soils* 43(1): 83-92.

[7] Dong ShouKun, Liu LiJun, Sun CongShu, Zhang Bing, Ma XiuFeng, Jiao GuangChun, Ma ChunMei, Gong ZhenPing. 2011. A study of the effects of nitrogen levels on nodule growth of soybean using ¹⁵N tracing. *Plant Nutrition and Fertilizer Science* 17 (4): 985-988.

[8] Gardezi, A.K., R. Ferrera-Cerrato, J. Kohashi, E.M. Engleman y M. Larqué-Saavedra. 1990. Potencial diferentes variedades de *Phaseolus vulgaris* de alta eficiencia en la fijación de nitrógeno en asociación con *Rhizobium leguminosarum* Biovar *Phaseoli*. *Agrociencia* 1(4): 25-39.

[9] Gardezi, A. K., I. D. Barceló-Quintal, V.M. Cetina-Alcalá, A. L. Bussy, M. U. Larqué-Saavedra, E. M. Saenz, A. Exebio-García, J. Pérez-Nieto, and M. A. Borja-Salin. 2005. Phytoremediation by *Leucaena leucocephala* in association with arbuscular endomycorrhiza and *Rhizobium* in soil polluted by Cr. *In* Callaos et al. The 9th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. VII: 289-298.

[10] Gardezi, A. K., E. Ojeda-Trejo, E. Mejia-Saenz, A. Exebio-García, U. Larque-Saavedra, S. Márquez-Berber, y V.M. Cetina –Alcalá. 2007. Effect of arbuscular endomycorrhizas associated with cow manure in *Leucaena leucocephala*. *In* Callaos et al. The 11th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. IV: 113-118.

[11] Gardezi, A. K., A. Exebio-García, E. Ojeda-Trejo, E. Mejia-Saenz, L. Tijerina-Chavez, M. Delgadillo-Piñon, U. Larqué-Saavedra, C. Villanueva-Verduzco, and H. Gardezi. 2008a. Growth response of *Leucaena leucocephala* associated with arbuscular endomycorrhizae to applications of organic matter in an irrigated soil with sewage water. *In* Callaos et al. The 12th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. IV: 153-158.

[12] Gardezi, A. K., E. Ojeda-Trejo, and S. R. Márquez-Berber. 2008b. Respuesta a la inoculación de *Glomus intraradix*, materia orgánica y dosis de fertilización fosfatada en el crecimiento de mezquite (*Prosopis* sp.). *Agroproductividad* 1(1): 24-28.

[13] Gardezi, A. K., I. D. Barceló, A. Exebio-García, E. Mejía-Saenz, U. Larqué-Saavedra, S. R. Márquez-Berber, C. Villanueva-Verduzco, H. Gardezi, and D. Talavera-Magaña. 2009. Cu²⁺ bioaccumulation by *Leucaena leucocephala* in symbiosis with *Glomus* spp.

and *Rhizobium* in copper-containing soil. In Callaos et al. The 13th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. II: 17-22.

[14] Gardezi, A. K., B. Figueroa-Sandoval, A. Exebio-García, S. R. Márquez-Berber, M. U. Larqué-Saavedra, E. Mejía-Saenz, D. Talavera-Magaña, M. E. Delgadillo-Piñon, C. Villanueva-Verduzco, and H. Gardezi. 2010. Effect of *Glomus intraradices* associated with different genotypes of *Phaseolus vulgaris* (common bean) in two soil types. In Callaos et al. Proceedings. The 14th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. II:5-10.

[15] Gardezi, A. K., S. R. Marquez-Berber, B. Figueroa-Sandoval, U. Larqué-Saavedra, M. Escalona-Maurice. 2012a. Endomycorrhizal inoculation effect on beans (*Phaseolus vulgaris* L.), oat (*Avena sativa* L.), and wheat (*Triticum aestivum* L.) growth cultivated in two soil types under greenhouse conditions. Journal of Systemics, Cybernetics and Informatics 10(5): 68-72.

[16] Gardezi, A. K., Márquez-Berber S. R., Figueroa-Sandoval, B., Iourii Nikolskii Gavrilov, Larqué-Saavedra U., Escalona-Maurice M. J. 2012b. Fungicide effect on *Glomus intraradices* in different genotypes of beans (*Phaseolus vulgaris* L.), oat (*Avena sativa* L.), and wheat (*Triticum aestivum* L.) growth cultivated in two soil types under greenhouse conditions. In Callaos et al. Proceedings. The 16th World Multiconference on Systemics, Cybernetics and Informatics. Orlando, Florida. U.S.A. I:61-66.

[17] George, E.; V. Romheld y H. Marschner. 1994. Contribution of mycorrhizal fungi to micronutrient uptake by plants. In J.A. Manthey, D.E. Crowley y D.G. Lewis (eds.). Biochemistry of metal micronutrients in the rhizosphere. London. pp. 93-109.

[18] Huang, W. 2009. Factors Contributing to the Recent Increase in U.S. Fertilizer Prices, 2002-08. Outlook AR-33. A Report from the Economic Research Service. USDA.

<http://www.ers.usda.gov/Publications/AR33/AR33.pdf>

[19] Irizar Garza, M. B.; Vargas Vázquez, P.; Garza García, D.; Tut y Couoh, C.; Rojas Martínez, I.; Trujillo Campos, A.; García Silva, R.; Aguirre Montoya, D.; Martínez González, J. C.; Alvarado Mendoza, S.; Grageda Cabrera, O.; Valero Garza, J.; Aguirre Medina, J. F. 2003. Respuesta de cultivos agrícolas a los biofertilizantes en la Región Central de México. Agricultura Técnica en México Vol. 29 No. 2 pp. 213-225.

[20] Lavelle P, and A. Spain. 2005. Soil Ecology. Springer Publisher, Dordrecht, The Netherlands.

[21] Novella L., R., A. M. Machado A., M. C. Ladrón de Guevara, y A. Vega B. 2003. Participación de los biofertilizantes en la producción del policultivo. Alimentaria 349: 133-135.

[22] Lindemann, W. C., and C. R. Glover. 2003. Nitrogen Fixation by Legumes. Guide A-129. Cooperative Extension Service. College of Agriculture and Home Economics. New Mexico State University.

[23] Plenchette, C., C. Clermont-Dauphin, J. M. Meynard, and J. A. Fortin. 2005. Managing arbuscular mycorrhizal fungi in cropping systems. Can J Plant Sci. 85:31-40.

[24] SIAP-SIACON. 2012. Information System for Agrifood Consultation. Database. Ministry of Agriculture. Mexico. www.siap.gob.mx

[25] Tawaraya, K. 2003. Arbuscular mycorrhizal dependency of different plant species and cultivars. Soil Science and Plant Nutrition 49 (5): 655-668.

[26] Tu, C.; Booker, F. L.; Watson, D. M.; Chen, X.; Ruffy, T. W.; Shi, W.; Hu, S. J. 2006. Mycorrhizal mediation of plant N acquisition and residue decomposition: impact of mineral N inputs. Global Change Biology. 12(5): 793-803.

[27] Yücel, C.; Özkan, H.; Ortas, I.; Yagbasanlar, T. 2009. Screening of wild emmer wheat accessions (*Triticum turgidum* subsp. *dicoccoides*) for mycorrhizal dependency. Turkish Journal of Agriculture and Forestry 33 (5): 513-523.



Dr Abdul Khalil Gardezi, paper main author.

Dr Abdul Khalil Gardezi is a distinguished scientist and academic member of the Hydro Science Center, Postgraduate College in Agriculture Science in Mexico, since 1981. He has received distinctions for teaching, research and service from 1988 until 2012. He has been selected for the originality of his research, presented as the best paper and oral presentation during 2003, 2005, 2008, 2009, 2010, 2011, and 2012 in international congresses in USA, England, Mexico and Australia. He has published more than 170 papers national and internationally. He has been honored among 2000 outstanding intellectuals of the 21st century by the International Biographical Center Cambridge, England.